

As a library, NLM provides access to scientific literature. Inclusion in an NLM database does not imply endorsement of, or agreement with, the contents by NLM or the National Institutes of Health.

Learn more: [PMC Disclaimer](#) | [PMC Copyright Notice](#)



[Atherosclerosis](#). Author manuscript; available in PMC 2020 Jun 7.

Published in final edited form as:

[Atherosclerosis](#). 2020 Mar; 296: 68–73.

Published online 2020 Jan 29. doi: [10.1016/j.atherosclerosis.2020.01.008](https://doi.org/10.1016/j.atherosclerosis.2020.01.008)

PMCID: PMC7276095

NIHMSID: NIHMS1586118

PMID: [32033778](#)

Calcium supplements: Good for the bone, bad for the heart? A systematic updated appraisal

[Marco B. Morelli](#), [Gaetano Santulli](#),* and [Jessica Gambardella](#)

Calcium is the most abundant mineral in the human body. Approximately 99% of the total body calcium is present, in form of hydroxyapatite – a lattice-like crystal composed of calcium, phosphorus, and hydroxide – in bones and teeth, where it essentially plays a structural role. The remaining calcium is in the extracellular fluid and in various tissues, where it regulates numerous processes, including vascular tone, muscle contraction-dilation, nerve impulse transmission, intracellular signaling, ATP production, and hormonal secretion.

The physiological concentration of calcium in the blood is tightly regulated ([Table 1](#)) within a range of 8.8–10.4 mg/dL by parathyroid hormone (parathormone, PTH), 1,25-dihydroxyvitamin D3 (active metabolite of vitamin D), and calcitonin (CT). In order to maintain a normal calcium level, and to avoid conditions of calcium deficiency, the National Institutes of Health and the Institute of Medicine recommend a daily calcium intake of 1000 mg for men aged 19–70 and women aged 19–50, to be increased up to 1200 mg for men aged 71 or older and women aged 51 or older.

Table 1

Normal values in adults.

| | Normal range (SI units) | Normal range (Conventional units) |
|--|-------------------------|-----------------------------------|
| Serum calcium | 2.2–2.6 mMol/L | 8.8–10.4 mg/dL |
| Ionized calcium ^a | 1.1–1.3 mMol/L | 4.4–5.2 mg/dL |
| PTH | 1.6–6.9 pMol/L | 15–65 pg/mL |
| Calcidiol, or 25-hydroxyvitamin D [25(OH)D] ^b | 50–200 nMol/L | 20–80 ng/mL |

PTH: parathyroid hormone; SI: International System.

^aAlbeit all calcium in the body is technically ionized, the term usually only applies to the free ionic fraction that is physiologically active in the blood.

^bIn contrast to 25(OH)D, circulating calcitriol, also known as 1,25(OH)2D, is generally not a good indicator of vitamin D status because it has a short half-life of 15 hours and serum concentrations are closely regulated by parathyroid hormone, calcium, and phosphate; levels of 1,25(OH)2D do not typically decrease until vitamin D deficiency is severe.

1. Physiology of calcium homeostasis

In addition to conferring strength to the skeleton, the deposition of calcium in the bone represents a source of the cation when its serum concentration decreases: indeed, free calcium can be released through osteoclastic bone resorption. An unbalanced equilibrium between bone calcium deposition and resorption affects the bone mineral density (BMD) resulting in skeletal fragility. The leading actor in the regulation of this equilibrium and calcium homeostasis is vitamin D, which allows the intestinal absorption of calcium, favoring its storage through bone calcium deposition. When the serum calcium content decreases, the sensory receptors on parathyroid glands promote PTH release, which activates bone resorption to restore blood calcium and promotes Vitamin D activation, to indirectly support the absorption of calcium from external sources. If PTH is the main stimulator of Vitamin D synthesis, Vitamin D, in turn, inhibits PTH secretion [1,2]. A normal BMD reflects the proper functioning of this elegant mechanism.

During the physiological bone growth, when the calcium demand increases, or in pathological conditions characterized by defective calcium absorption, a higher calcium intake could be necessary to preserve BMD. However, the positive effects of calcium supplementation on skeleton fragility seem to be in contrast with the risk of side effects, such as cardiovascular events.

2. Impact of calcium supplements on the cardiovascular system

Although a balanced dietary intake of calcium and vitamin D is critical for our body in order to regulate critical mechanisms, the role of calcium and vitamin D supplementation in adults and elderly people remains not fully understood. In this issue of Atherosclerosis, Hulbert and colleagues show that calcium supplementation, but not dietary calcium, positively correlates with abdominal aorta calcification in postmenopausal women [3], suggesting that vascular calcification might contribute to the cardiovascular events observed in calcium supplement users. Several prospective cohort and randomized studies have demonstrated that an adequate calcium intake is associated with beneficial cardiovascular effects, including protection from hypertension, vascular disease, and stroke; moreover, calcium in association with vitamin D seems to be protective against coronary heart disease (CHD).

Based on these perceptions about calcium as a key component for the bone structure and critical for fundamental cellular processes, during the last decades we have observed a huge diffusion of calcium supplements and food enriched in calcium. Indeed, more than 50% of elderly men and almost 70% of elderly women in the US are considered regular users of calcium supplements.

3. Effects of calcium supplementation on BMD: results from meta-analyses

Several clinical trials have tested the efficacy of calcium supplementation in children and adolescents, overall showing a positive effect of calcium introduction on BMD [4]. Instead, it is not clear whether these effects are also detectable in the mature skeleton. One randomized clinical trial conducted in 354 females in the pubertal stage, examined the long-time effects of calcium supplement on total-body BMD, from child to young adult [5]; the study revealed an increased BMD in the supplemented group at year 4, but this outcome vanished thereafter, supporting the hypothesis that calcium supplementation effects vary over time: calcium supplementation appears to influence bone accretion during pubertal growth spurt, losing most of the effects from young adult onwards.

In Western countries, the average calcium intake by diet is around 700–900 mg/day, which could be not enough to guarantee a proper BMD, especially in the risk population. Therefore, calcium supplementation has been proposed also for adult people and for the aging population to prevent or treat the pathological reduction of BMD that can lead to osteoporosis. To test the actual efficacy of this approach in preserving the BMD, various studies have been conducted and some meta-analyses well summarize the issue [6,7]. The first meta-analysis considered trials examining the effect of calcium supplementation on BMD and the risk of fractures in postmenopausal women showing a weak positive effect of calcium supplementation on BMD, with an increase of 2,05% of total body bone density after two or more years of treatment, respect to baseline [6]. This increase seems to have a rather low clinical relevance as it was associated with a reduced trend of the risk of experiencing only vertebral fractures (the risk was not reduced when evaluating non-vertebral fractures). Another meta-analysis considered the general population (men and women) aged over 50, including a total of 51 randomized control trials focused on calcium supplementation and changes in BMD [7]; overall, the analysis showed that calcium supplementation increases BMD by a maximum of 1,8%, starting from the first year of treatment without a progressive increase over time. Interestingly, this study indicates that the increase in BMD in response to calcium supplementation is of equal entity to what detected in response to calcium intake from diet. A meta-analysis that examined 33 randomized trials involving a total of 51,145 participants, demonstrated that the use of supplements that included calcium, vitamin D, or both compared with placebo or no treatment was not associated with a lower risk of fractures among community-dwelling older adults [8]. The most recent meta-analysis, as


sessing 6 randomized-controlled trials of vitamin D combined with calcium vs placebo or no treatment shows that neither intermittent nor daily dosing with standard doses of vitamin D alone is associated with reduced risk of fracture; however, combined treatment with both vitamin D and calcium is associated with a 16% (95% CI, 3%–28%) reduction in the risk of hip fracture [9].

4. Calcium and vitamin D: friends or foes?

As mentioned above, calcium supplementation alone seems to have a minimum clinical impact in terms of risk of fractures. This outcome is not surprising if we consider that, especially in aged populations, a key issue is calcium absorption, due to a defective vitamin D bioavailability and/or activity. Indeed, vitamin D is the key regulator of intestinal calcium absorption; with aging, an increase of the enzyme involved in vitamin D catabolism occurs, alongside with a vitamin D resistance. Therefore, if a main problem is in vitamin D-dependent calcium absorption, the increase of calcium intake by diet or supplementation could not be adequate to overcome the insufficient intestinal absorption. In light of this consideration, the addition of vitamin D to calcium administration should be more effective. Indeed, the effects of calcium supplementation on the risk of fractures is amplified by co-administration of vitamin D [10], particularly for hip fractures. In general, combining calcium and Vitamin D reduces by 10–20% the occurrence of non vertebral fractures [11]. Nevertheless, a recent trial has shown that daily supplementation with high-dose vitamin D does not result in a lower incidence of cardiovascular events or invasive cancer than placebo [12].

5. Calcium and cardiovascular risk: controversy even in meta-analyses; do we need more cowbell?

Most recently, some investigators started raising their eyebrows on this topic, because of the potential detrimental effects of high calcium intake (with or without vitamin D) on cardiovascular risk [13]. One of the principal mechanisms triggering major concerns in some researchers is the acutely increased serum calcium levels observed after calcium supplement intake; in fact, calcium intake from dietary sources does not seem to increase cardiovascular risk, while calcium supplements might raise CHD risk (Fig. 1).



[Fig. 1.](#)

Calcium supplements rapidly increase circulating calcium and high levels of circulating calcium have been shown to increase cardiovascular risk; one of the possible mechanisms is the progressive ectopic calcification of the arteries, which could lead to coronary artery disease. On the other hand, dietary calcium intake has been shown to provide calcium via a slower assimilation, thereby allowing to maintain over time physiological calcium levels, a condition that seems to reduce the risk of cardiovascular disease.

Large observational studies have shown that augmented concentrations of serum calcium can increase the risk of myocardial infarction (MI), therefore it seemed logical to consider potentially dangerous to rise peaks of circulating calcium since this process can advance dangerous (ectopic) calcifications. Reid and collaborators reported that ionized calcium concentration increased from a mean of 1.22–1.30 mmol/L following supplementation with 1 g of calcium, either as citrate or as lactate gluconate [13]. Therefore, it is plausible that augmented plasma levels of calcium might alter the systemic balance in favor of calcification [13], which becomes a serious health risk when it occurs as ectopic calcification. The controversy on the actual role of calcium supplementation on cardiovascular risk has not been solved by meta-analyses (Table 2) of published studies, which have generated inconsistent results [14–21], showing how this important topic is debated and exposing the current uncertainty about optimal doses and regimens for supplementation and their overall effectiveness.

Table 2

Meta-analyses assessing the association between DCSs or CSs intake (with or without Vitamin D) and cardiovascular events.

| Authors, Year | Meta-analysis for CHD, MI, CVD, and all-cause mortality | Total number of subjects | Ethnicity – Sex (age) | Follow –up (average or range of years) | Outcome | Ref. |
|----------------------|--|---|--|---|---|--------------------|
| Bolland et al., 2010 | 15 studies | 8151 (in 15 trials eligible for inclusion); plus 11,921 participants (in 11 trial level data) | Different ethnicity; males and females (from 51 to 77 years old) | Average of 3.6 years (15 trials for inclusion) and 4 years (11 trials level data) | In five studies contributing patient level data (5 studies), 143 participants allocated to calcium had an MI compared with 111 allocated to placebo (hazard ratio 1.31, 95% confidence interval 1.02 to 1.67, $p = 0.035$); composite endpoint of MI, stroke, or sudden death (hazard ratio 1.18, 95% confidence interval 1.00 to 1.39, $p = 0.057$), and death (1.09, 0.96 to 1.23, $p = 0.18$). The meta-analysis of trial level data (11 studies) yielded similar results: 296 participants had a myocardial infarction (166 allocated to calcium, 130 to placebo), with an increased incidence of myocardial infarction in those allocated to calcium (pooled relative risk 1.27, 95% confidence interval 1.01 to 1.59, $p = 0.038$). | 14 |
| Bolland et al., 2011 | 2 trials of co-administered calcium and vitamin D with CVD outcome data (WHI CaD | 28,072 | Different ethnicity; only females (post-menopausal women) | Average 5.9 years | CSs with or without vitamin D modestly increase the risk of CVD events, especially MI. In | 15 |

Acknowledgments

Financial support

The Santulli's Lab is supported by the National Institutes of Health (R01-HL146691, R01-DK123259, R00-DK107895, P30-DK020541, and R01-DK033823 to Prof. Gaetano Santulli) and by the American Heart Association (AHA-20POST35211151 to Dr. Jessica Gambardella).

Footnotes

Declaration of competing interest

The authors declared that they do not have anything to disclose regarding conflict of interest with respect to this manuscript.

Contributor Information

Marco B. Morelli, Department of Medicine, Wilf Family Cardiovascular Research Institute, Albert Einstein College of Medicine, New York, NY, USA.

Gaetano Santulli, Department of Medicine, Wilf Family Cardiovascular Research Institute, Albert Einstein College of Medicine, New York, NY, USA. Department of Molecular Pharmacology, Einstein-Mount Sinai Diabetes, Research Center (ES-DRC), The "Norman Fleischer" Institute for Diabetes and Metabolism, Albert Einstein College of Medicine, New York, NY, USA. International Translational Research and Medical Education Consortium (ITME), Naples, Italy. Department of Advanced Biomedical Sciences, "Federico II" University, Naples, Italy.

Jessica Gambardella, Department of Medicine, Wilf Family Cardiovascular Research Institute, Albert Einstein College of Medicine, New York, NY, USA. International Translational Research and Medical Education Consortium (ITME), Naples, Italy.

References

[1] Gambardella J, De Rosa M, Sorriento D, et al., Parathyroid hormone causes endothelial dysfunction by inducing mitochondrial ROS and specific oxidative signal transduction modifications, *Oxid. Med. Cell. Longev* 2018 (2018) 9582319. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

- [2] Pascale AV, Finelli R, et al., Vitamin D, parathyroid hormone and cardiovascular risk: the good, the bad and the ugly, *J. Cardiovasc. Med* 19 (2018) 62–66. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
- [3] Hulbert M, Turner ME, Hopman WM, Changes in vascular calcification and bone mineral density in calcium supplement users from the Canadian multi-center osteoporosis study (CaMOS), *Atherosclerosis* 296 (2020) 83–90. [[PubMed](#)] [[Google Scholar](#)]
- [4] Johnston CC Jr, Miller JZ, Slemenda CW, et al., Calcium supplementation and increases in bone mineral density in children, *N. Engl. J. Med* 327 (1992) 82–87. [[PubMed](#)] [[Google Scholar](#)]
- [5] Matkovic V, Goel PK, Badenhop-Stevens NE, et al., Calcium supplementation and bone mineral density in females from childhood to young adulthood: a randomized controlled trial, *Am. J. Clin. Nutr* 81 (2005) 175–188. [[PubMed](#)] [[Google Scholar](#)]
- [6] Shea B, Wells G, Cranney A, et al., Meta-analyses of therapies for postmenopausal osteoporosis. VII. Meta-analysis of calcium supplementation for the prevention of postmenopausal osteoporosis, *Endocr. Rev* 23 (2002) 552–559. [[PubMed](#)] [[Google Scholar](#)]
- [7] Tai V, Leung W, Grey A, et al., Calcium intake and bone mineral density: systematic review and meta-analysis, *BMJ* 351 (2015) h4183. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
- [8] Zhao JG, Zeng XT, Wang J, et al., Association between calcium or vitamin D supplementation and fracture incidence in community-dwelling older adults: a systematic review and meta-analysis, *J. Am. Med. Assoc* 318 (2017) 2466–2482. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
- [9] Yao P, Bennett D, Mafham M, et al., Vitamin D and calcium for the prevention of fracture: a systematic review and meta-analysis, *JAMA Netw. Open* 2 (2019) e1917789. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
- [10] Tang BM, Eslick GD, Nowson C, et al., Use of calcium or calcium in combination with vitamin D supplementation to prevent fractures and bone loss in people aged 50 years and older: a meta-analysis, *Lancet* 370 (2007) 657–666. [[PubMed](#)] [[Google Scholar](#)]
- [11] Avenell A, Gillespie WJ, Gillespie LD, et al., Vitamin D and vitamin D analogues for preventing fractures associated with involutional and post-menopausal osteoporosis, *Cochrane Database Syst. Rev* (2009) CD000227. [[PubMed](#)]
- [12] Manson JE, Cook NR, Lee IM, et al., Vitamin D supplements and prevention of cancer and cardiovascular disease, *N. Engl. J. Med* 380 (2019) 33–44. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
- [13] Reid IR, Bolland MJ, Grey A, Does calcium supplementation increase cardiovascular risk? *Clin. Endocrinol* 73 (2010) 689–695. [[PubMed](#)] [[Google Scholar](#)]
- [14] Bolland MJ, Avenell A, Baron JA, et al., Effect of calcium supplements on risk of myocardial infarction and cardiovascular events: meta-analysis, *BMJ* 341 (2010) c3691. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
- [15] Bolland MJ, Grey A, Avenell A, et al., Calcium supplements with or without vitamin D and risk of cardiovascular events: reanalysis of the Women’s Health Initiative limited access dataset and meta-analysis, *BMJ* 342 (2011) d2040. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
- [16] Rejnmark L, Avenell A, Masud T, et al., Vitamin D with calcium reduces mortality: patient level pooled analysis of 70,528 patients from eight major vitamin D trials, *J. Clin. Endocrinol. Metab* 97 (2012) 2670–2681. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
- [17] Wang X, Chen H, Ouyang Y, et al., Dietary calcium intake and mortality risk from cardiovascular disease and all causes: a meta-analysis of prospective cohort studies, *BMC Med* 12 (2014) 158. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

[18] Asemi Z, Saneei P, Sabihi SS, et al., Total, dietary, and supplemental calcium intake and mortality from all-causes, cardiovascular disease, and cancer: a meta-analysis of observational studies, *Nutr. Metabol. Cardiovasc. Dis* 25 (2015) 623–634. [[PubMed](#)] [[Google Scholar](#)]

[19] Lewis JR, Radavelli-Bagatini S, Rejnmark L, et al., The effects of calcium supplementation on verified coronary heart disease hospitalization and death in postmenopausal women: a collaborative meta-analysis of randomized controlled trials, *J. Bone Miner. Res* 30 (2015) 165–175. [[PubMed](#)] [[Google Scholar](#)]

[20] Chung M, Tang AM, Fu Z, et al., Calcium intake and cardiovascular disease risk: an updated systematic review and meta-analysis, *Ann. Intern. Med* 165 (2016) 856–866. [[PubMed](#)] [[Google Scholar](#)]

[21] Yang C, Shi X, Xia H, et al., The evidence and controversy between dietary calcium intake and calcium supplementation and the risk of cardiovascular disease: a systematic review and meta-analysis of cohort studies and randomized controlled trials, *J. Am. Coll. Nutr* (2019) 1–19. [[PubMed](#)]