

- Fasting glucose 70–95 mg/dL (3.9–5.3 mmol/L) and either
- One-hour postprandial glucose 110–140 mg/dL (6.1–7.8 mmol/L) or
- Two-hour postprandial glucose 100–120 mg/dL (5.6–6.7 mmol/L)

GLYCEMIC GOALS IN PREGNANCY

Recommendations

15.7 Fasting, preprandial, and postprandial blood glucose monitoring are recommended in individuals with diabetes in pregnancy to achieve optimal glucose levels. Glucose goals are fasting plasma glucose <95 mg/dL (<5.3 mmol/L) and either 1-h postprandial glucose <140 mg/dL (<7.8 mmol/L) or 2-h postprandial glucose <120 mg/dL (<6.7 mmol/L). **B**

term safety data. **E**

15.16 Metformin, when used to treat polycystic ovary syndrome and induce ovulation, should be discontinued by the end of the first trimester. **A**

15.8 Due to increased red blood cell turnover, A1C is slightly lower during pregnancy in people with and without diabetes. Ideally, the A1C goal in pregnancy is <6% (<42 mmol/mol) if this can be achieved without significant hypoglycemia, but the goal may be relaxed to <7% (<53 mmol/mol) if necessary to prevent hypoglycemia. **B**